

RESEARCH

Open Access



Suicidal behaviors in the entertainment industry: a preliminary exploration of the interplay between work scheduling, social support, and wellbeing in Australia

Daniel Zarate¹, Christopher Sonn¹, Adrian Fisher¹ and Vasileios Stavropoulos^{1,2*}

Abstract

Objective: Workers of the Australian entertainment industry exhibit disproportionately high rates of impaired psychological wellbeing and suicidal behaviors, with such rates being exacerbated by the negative impact of working long and odd hours (Work Scheduling Impact; WSI). Nonetheless, stable and secure social support networks may buffer the risks associated with such systemic difficulties.

Methods: The responses of 1302 Australian entertainment industry workers (50.3% females, M_{age} 38.39 years) on the Multidimensional Scale of Perceived Social Support, the Short Form Health Survey, WSI, and suicidal behaviors questions were examined via moderation analyses.

Results: Higher social support and lower WSI appeared to reduce the suicidal ideation of those experiencing poorer mental health, while lower WSI further enhanced social support's positive effect.

Conclusions: Findings highlight the likely detrimental effect of WSI regarding the suicidal ideation reported by vulnerable Australian entertainment industry workers and stress the importance of the social support they experience.

Public health implications: Interventions attempting to increase social support could improve inherent conditions associated with the Australian entertainment industry. Similarly, the negative effect of working long and odd hours on workers' mental health and suicidal behaviors indicates the need to regulate the industry appropriately.

Keywords: Australian entertainment industry, Mental health, Multidimensional perceived social support, Suicide, Moderation analyses

Introduction

Workers of the entertainment industry experience low rates of mental wellbeing and suicidal behaviors [1–3]. Difficulties associated with the nature of the entertainment industry, such as unstable employment and a 'toxic culture' have been proposed to underpin poorer mental

wellbeing [2, 4]. For example, Van Rens and Heritage [3] reported that despite having high levels of resilience, circus performers showed higher depression, anxiety, and stress than 88%, 86%, and 77% of adults, respectively. Moreover, Van Den Eynde and colleagues [2] revealed that 44% of workers of the Australian entertainment industry have moderate to severe anxiety, 15% moderate to severe depression, and more importantly, four to five times greater incidence of suicide planning, resulting in more than double attempts compared to other Australians. Additionally, their work scheduling impact (WSI)

*Correspondence: Vasileios.stavropoulos@vu.edu.au

¹ Institute for Health and Sport, Victoria University, Melbourne, Australia
Full list of author information is available at the end of the article



due to working long and odd hours disrupts individuals' social lives, could adversely affect their health, sleep [5], overall wellbeing [6–8], and likely increase their suicide risk. Thus, it is plausible that the WSI may associate with entertainment industry workers' increased suicide behaviors and mental health issues.

In the present study, such suicide behaviors are considered to involve ideation, planning, and attempts [9–11]. Ideation is defined as the contemplations, wishes, and preoccupations with death and suicide; planning is the precursor of potentially lethal attempts; while attempts involve self-inflicted behaviors with non-fatal outcomes [12, 13]. Interestingly, factors related to suicidal behaviors may differently influence suicidal ideation, planning, or attempts over time [14]. For example, Joiner [15] illustrates that while mental disorders may result in higher suicide ideation, only certain disorders, such as posttraumatic stress disorder, may increase the risk for suicide attempts. Moreover, Klonsky and colleagues [14] suggest longitudinal variations of the same factors, such as loneliness and isolation, due to cumulative effects, with ideation being the initial step and planning and attempts following. To address such issues, the current study integrates four different measures concerning one's suicidal behaviors, including ideation in the last 12 months and over one's life, planning, and attempts. While assessing these distinct yet interrelated aspects, the present research considers self-reported social support as a potential protective condition [2, 16].

Buffering effect of social support networks

Social support refers to one's perception of social integration, bonds, and closeness to others in their surroundings [17]. Pearson [17] highlights three elements present in one's sense of social support directly related to psychological wellbeing: (i) being cared for and loved, (ii) being valued, and (iii) experiencing a sense of belonging to a social group. Stable social support provides basic psychological and developmental needs such as belonging, self-esteem, and security with a direct (and positive) impact on depression, emotional stability, and suicidal behaviors [18]. Additionally, scholars proposed that social support has a buffering effect on burnout and work-related stress, as it may help individuals relax/distress from their perceived work demands [19]. Accordingly, one could assume that a strong social support network could buffer/moderate the risk of poor mental health and adverse WSI on suicidal behaviors among those working in entertainment.

Current research

Systemic difficulties associated with the entertainment industry, such as WSI, have compromised workers'

wellbeing [2, 3]. Similarly, the buffering effect that social support systems have on daily stressors and poorer mental health has been demonstrated [19]. Despite this, there is limited research regarding the significance of the interplay between WSI (as a risk factor) and social support (as a potential buffer for suicidal behaviors) in workers of the Australian entertainment industry. Additionally, complex issues continue to affect the Australian entertainment industry (for example, a recent report revealed significant issues in the industry associated with toxicity, financial instability, and low mental wellbeing [20]), suggesting the need to explore these issues further.


The present study aims to address this gap by emphasizing how the WSI-Social support interaction could associate with the likelihood of Australian, entertainment industry workers experiencing lower wellbeing, to engage in suicidal behaviors. Specifically, it combines unique conceptual and methodological strengths to translate its findings into specific practical implications. Findings aspire to provide vital information to policymakers via: (1) shedding light on the need, and the directions, of regulating the precarious position of those working in the entertainment industry to potentially regulate their WSI; and (2) demonstrating the importance of social support networks in suicide prevention. To address these aims, the following hypotheses were formulated:

- Workers experiencing lower mental health/wellbeing and higher WSI will report higher suicidal ideation in the last 12 months (H_1), over their lifetime (H_2), suicide planning (H_3), and suicide attempts (H_4 , Fig. 1 panel A) compared to those showing lower WSI.
- Workers experiencing lower mental health/wellbeing and higher levels of social support will report lower suicidal ideation in the last 12 months (H_5), suicidal ideation in participants' lifetime (H_6), suicide planning (H_7), and suicide attempts (H_8 , Fig. 1 panel B) compared to those showing lower social support.
- Workers experiencing higher levels of social support will report lower suicidal ideation in the last 12 months (H_9), suicidal ideation in participants' lifetime (H_{10}), suicide planning (H_{11}), and suicide attempts (H_{12} , Fig. 1 panel C and D), compared to their counterparts experiencing concurrently the same level of mental health/wellbeing and WSI adversity.

Method

Participants

A total of 2409 workers of the Australian entertainment industry completed an online survey for a project titled "Working in the Australian Entertainment Industry: Final



Report” between 23/01/2015 and 08/05/2015 [2]. The original survey included a series of questionnaires beyond the scope of the current study. A filter question/item related to one’s experience of suicidal thoughts and/or actions, which constitutes the specific interest of the present study, was positively answered by 1302 respondents, who additionally addressed the suicide-specific measures analyzed here. Thus, the current study used a sample of 1302 participants ($M_{age} = 38.39, SD_{age} = 0.98$; 50.3% females). Missing responses ($n = 331, 6.1\%$) were completely at random (MCAR Little’s test: $\chi^2_{[1096]} = 0.065$), and thus a listwise deletion was conducted. A-priori power analyses via the G-power software [21] indicated that for two tails of significance, $F^2 = 0.15$, α error probability = 0.05, Power ($1 - \beta$ error probability) = 0.95 and seven predictors (i.e., wellbeing/mental health, WSI, social support, wellbeing/mental health x WSI, wellbeing/mental health x social support, social support x WSI, wellbeing/mental health x social support x WSI), a total sample of 89 was required for an actual power of 0.95, at a critical $t = 1.989, df = 81$. Workers of this industry

belonged to several employment categories ranging from musicians, circus performers, dancers, managers, directors, and sound technicians. Supplementary Table 1 presents participants’ key demographic statistics.

Measures

The *Multidimensional Scale of Perceived Social Support (MSPSS)* was employed to measure subjective perceived social support [22]. A total of 12 items loads on three subscales (four per subscale, i.e., friends, family, and significant other) rated on a seven point-Likert scale (1 = *very strongly disagree* to 7 = *very strongly agree*) with lower scores indicating lower perceived support. Examples of items include “*There is a special person who is around when I am in need*”. Total item responses were added to reflect one’s social support (as a whole and per dimension), ranging between 12–84, and 4–48, respectively. The internal consistency for one’s overall social support in the present study was satisfactory (Cronbach’s $\alpha = 0.91$, McDonald’s $\omega = 0.91$, see Supplementary Table 2 [23]).

The *Short Form 12 items Health Survey (SF12)* was used to assess mental health [24]. A total of 12 items loads on two subscales (i.e., mental, and physical component) and include items such as “*As a result of emotional problems, have you accomplished less than you would like?*”. Standardised scores suggest a mean of 50 ($SD=10$) with lower scores indicating lower mental wellbeing. The internal consistency for the present study was satisfactory ($\alpha=0.88$; $\omega=0.88$; Supplementary Table 2). Considering the focus of the study, we only employed the mental health component (MHC) of the scale.

Work Schedule Impact (WSI)

The scale determines the impact on wellbeing that a particular work schedule has [2]. The scale consists of five items loading on two subscales (intra-individual and inter-individual impact) and includes items such as “*I have trouble maintaining my social life as a result of my work schedule*”. Items were rated on a five point-Likert scale with higher scores indicating a higher impact due to work schedules. The questionnaire exhibited internal consistency ($\alpha=0.81$; $\omega=0.81$; see Supplementary Table 2) and appropriate factorizable properties (EFA; $KMO=0.76$, Bartlett’s χ^2 [15] = 2861; $p<0.001$), item loading 0.50 to 0.92; see Supplementary Table 3).

Suicidal behaviors

Following previous suggestions [2, 9], specific questions with dichotomous outcomes (yes/no) were employed to assess all three aspects of suicidal behaviors (i.e., ideation, planning, and attempts). Three questions were employed to assess participants’ suicide ideation in their lifetime (i.e., “*In your lifetime, have you ever felt that life was not worth living?*”) and three questions to assess participants’ suicide ideation in the last 12 months (i.e., “*In the last 12 months, have you ever thought of taking your life, even if you would not really do it?*”). Scores were added for suicidal ideation in the last 12 months and lifetime, awarding one point per each ‘yes’ with a score range of 0–3 (high scores represent high suicidal ideation). These questions showed excellent internal consistency ($\alpha=0.87$; $\omega=0.87$; see Supplementary Table 2). Similarly, the question “*Have you planned to complete suicide?*” was employed to assess participants’ intention to act on suicidal thoughts without necessarily involving preparatory behaviors to die by suicide. Finally, the question “*Have you ever made an attempt to take your life?*” was employed to assess participants’ suicide attempts.

Procedure

A Victoria University research team, in conjunction with Entertainment Assist, conducted a project titled

‘Working in the Australian Entertainment Industry: Final Report’ attempting to identify health and wellbeing concerns for those who work in this industry [2] was approved by the Victoria University Ethics Committee (HRE14-270). Eligible participants, adult workers of the Australian creative industry, were invited to complete an online survey advertised through Entertainment Assist’s membership email list and their Facebook site. Before completing the survey, participants accessed information about the voluntary and anonymous nature of the study, its aims, significance, and their right to withdraw at any point and without repercussions via the Plain Language Information Statement.

Statistical analysis

A series of regression-based moderation analyses were conducted on IBM-SPSS 26 using the Process macro [25]. Specifically, this study used a model with two moderators (WSI and MSPSS) aiming to assess (i) the rate of change in the relationship between mental health (SF12) and suicidal behaviors at different levels of these moderators and (ii) the concurrent associations of both moderators. The moderating association of WSI on the relationship between mental health and suicidal behaviors was assessed by the interaction $SF12*WSI$, and the moderating association of MSPSS on the relationship between mental health and suicidal behaviors was assessed by the interaction $WSI*MSPSS$. Finally, a higher order interaction between the independent variable and both moderators ($SF12*WSI*MSPSS$) was used to observe the concurrent association of the interplay between work schedule and social support on the relationship between mental health and suicidal behaviors (for visual clarification see Fig. 1 panel C and D). Following suggestions outlined in Hayes [25], mean centring was conducted on variables that defined products, and simple slope analysis was used to visualize conditioning values at the 16th, 50th, and 84th percentiles. Subsequently, the Johnson-Neyman technique was used to provide more detail in visualizing moderating effects [25].

Several models were employed to test the hypothesized relationships. To test H_1 , H_5 , and H_9 , SF12 was used as the independent variable, WSI, MSPSS and their interactions were the moderators, and suicidal ideation in the last 12 months the outcome variable. The same model/structure was then reapplied with different outcome variables: (a) the suicidal ideation in participants’ lifetime as the outcome was employed to address H_2 , H_6 , and H_{10} ; (b) with suicide planning as the outcome regarding H_3 , H_7 , and H_{11} ; and (c) suicide attempts as the outcome variable considering H_4 , H_8 , and H_{12} . Due to the last two models referring to binary outcomes, logistic regression

instead of linear was calculated, while bootstrapping at 1000 resamples was applied for all analyses.

Results

To assess the moderating association of the interplay between work scheduling (WSI) and social support (MSPSS) on the relationship between mental wellbeing and suicidal ideation over the last 12 months (hypotheses H₁, H₅, and H₉), SF12 was employed as the independent variable, WSI as the moderator, MSPSS as the moderating moderator, and suicidal ideation in the last 12 months as the outcome variable. The model explained 27.3% of the variance in suicidal ideation during participants' last 12 months ($R^2=0.2729$, $F_{(7, 1294)}=69.380$, $p<0.001$), with SF12 ($b_{11}=-0.042$, $p<0.001$), WSI ($b_{12}=0.034$, $p<0.001$), and MSPSS ($b_{13}=-0.372$, $p<0.001$) as significant predictors.

Considering H₁, the moderating role of WSI on the relationship between SF12 and suicidal ideation in the last 12 months ($b_{14}=-0.001$, $p=0.724$) was not significant. Considering H₅, the moderating role of MSPSS on the relationship between SF12 and suicidal ideation in the last 12 months ($b_{15}=0.010$, $p=0.008$) was significant. As predicted, when MSPSS increased by one unit, the coefficient for the moderating effect of SF12 on suicidal ideation in the last 12 months increased by 0.01 (Table 1 and Fig. 2-left panel). Finally, considering H₉, the three-way interaction SF*WSI*MSPSS was significant ($b_{16}=-0.001$, $p=0.018$) explaining 0.32% of the

variance. When MSPSS increased by one unit, the coefficient for the moderating effect of SF12 and WSI on suicidal ideation in the last 12 months decreased by 0.001 (see Fig. 2, right panel for low, medium and high levels of MSPSS). Interestingly, the interaction between SF12 and WSI only became significant at MSPSS 85th percentile (MSPSS > 9.78; see Supplementary Table 4). In other words, the exacerbating role of WSI on the relationship between mental health and suicide ideation in the last 12 months was significantly mitigated at high levels of perceived social support.



To assess the moderating effects of role of the interplay between work scheduling (WSI) and social support (MSPSS) on the relationship between mental wellbeing and participants' suicide ideation over their lifetime (hypotheses H₂, H₆, and H₁₀), WSI was used as moderator, MSPSS as moderating moderator, and suicidal ideation in participants' lifetime as the outcome. The model explained 18.3% of the variance ($R^2=0.1824$, $F_{(7, 1294)}=41.251$, $p<0.001$), with SF12 ($b_{21}=-0.025$, $p<0.001$), WSI ($b_{22}=0.029$, $p<0.001$), and MSPSS ($b_{23}=-0.305$, $p<0.001$) being significantly associated to suicidal ideation in a participants' lifetime.

Considering H₂, the moderating role of WSI on the relationship between SF12 and suicide ideation in participants' lifetime ($b_{24}=0.002$, $p=0.002$) was significant. When WSI increased by one unit, the relationship between SF12 and suicidal ideation in participants' lifetime increased by 0.002 (Table 1 and Fig. 3, left panel).

Table 1 Estimating regression coefficients with 'Suicidal ideation in the last 12 months' and 'Suicidal ideation in participants lifetime' as outcome variables

Variable	<i>b</i>	BSE	<i>t</i>	<i>p</i>	BLLCI	BULCI
Outcome Variable—Suicidal Ideation in the last 12 months						
<i>α</i> : Constant/Intercept	1.0023	.0347	30.0849	.0001	.9325	1.0686
<i>b</i> ₁₁ : SF12	-.0418	.0033	-12.4114	.0001	-.0484	-.0356
<i>b</i> ₁₂ : WSI	.0341	.0078	4.5814	.0001	.0186	.0494
<i>b</i> ₁₃ : MSPSS	-.3720	.0420	-8.6329	.0001	-.4530	-.2873
<i>b</i> ₁₄ : SF12*WSI	-.0002	.0005	-.3528	.7243	-.0013	.0008
<i>b</i> ₁₅ : SF12*MSPSS	.0103	.0033	2.6408	.0084	.0037	.0167
<i>b</i> ₁₆ : SF12*WSI*MSPSS	-.0014	.0004	-2.3729	.0178	-.0022	-.0006
Outcome Variable—Suicidal Ideation in their lifetime						
<i>α</i> : Constant/Intercept	1.9618	.0351	57.3726	.0001	1.8930	2.0311
<i>b</i> ₂₁ : SF12	-.0253	.0035	-7.3337	.0001	-.0321	-.0185
<i>b</i> ₂₂ : WSI	.0295	.0079	3.8673	.0001	.0134	.0447
<i>b</i> ₂₃ : MSPSS	-.3052	.0448	-6.9498	.0001	-.3923	-.2172
<i>b</i> ₂₄ : SF12*WSI	.0019	.0006	3.0794	.0020	.0007	.0030
<i>b</i> ₂₅ : SF12*MSPSS	-.0036	.0037	-.8903	.3735	-.0109	.0036
<i>b</i> ₂₆ : SF12*WSI*MSPSS	-.0002	.0005	-.4089	.6827	-.0011	.0007

b Bootstrapped estimated value of unstandardised regression coefficient, *BSE* Bootstrapped standard error, *t* T-test statistic, *p* Probability, *BLLCI* Bootstrapped lower-level confidence interval, *BULCI* Bootstrapped upper-level confidence interval, *SF12* Short Form 12 items Mental Health Component, *WSI*/Work Schedule Impact, *MSPSS* Multidimensional Scale of Perceived Social Support



Considering H_6 and H_{10} , the moderating role of MSPSS on the relationship between SF12 and suicidal ideation in the last 12 months ($b_{25} = -0.004, p = 0.373$) and the three-way interaction $SF12*WSI*MSPSS$ ($b_{26} = -0.001, p = 0.683$) were not significant (see Fig. 3, right panel regarding low, medium and high MSPSS).

To assess the moderating role of the interplay between work scheduling (WSI) and social support (MSPSS) on the relationship between mental wellbeing and suicide planning (hypotheses H_3, H_7, H_{11}), a logistic regression model with SF12, WSI, and MSPSS as independent variables and suicide planning as the outcome variable was used. Interaction terms between independent variables were included in the model to assess potential moderating associations. The model represented a significant improvement with respect to a null model ($\chi^2_{[6]} = 77.149, p < 0.001$) and correctly classified 70.3% of cases. The Nagelkerke pseudo R^2 indicated that the predictors

accounted for approximately 8.4% of the variance in suicide planning. However, no single predictor or interaction term significantly predicted suicide planning (Table 2). Given that Wald's coefficient was largest for $SF12*WSI$ ($b_{34} = 0.340$) and $SF12*MSPSS$ ($b_{35} = 0.370$), it is likely that the interaction terms account for most of the overlapping variance otherwise associated with single predictors.

Finally, to assess the moderating role of the interplay between work scheduling (WSI) and social support (MSPSS) on the relationship between mental wellbeing and suicide attempts (hypotheses H_4, H_8, H_{12}), a logistic regression model was run with SF12, WSI, and MSPSS as independent variables and suicide attempts as the outcome variable. Interaction between independent variables was also included in the model to assess potential moderating effects. The model represented a significant improvement with respect to a null model

($\chi^2_{[6]} = 38.331, p < 0.001$) and correctly classified 86.5% of cases. The Nagelkerke pseudo R^2 indicated that the predictors accounted for approximately 5.5% of the variance. However, as with suicide planning, no single variable or interaction term was significantly associated with suicide planning (Table 2). In this case, larger Wald coefficients were observed for the three-way interaction SF12*WSI*MSPSS ($b_{46} = 0.305$), MSPSS ($b_{43} = 0.264$), and SF12 ($b_{41} = 0.258$), suggesting their comparative importance.

Discussion

Moderating effect of work scheduling impact (WSI) on the relationship between mental health and suicidal behaviors

This study has been (to the best of the authors’ knowledge) the first to distinctly evaluate the buffering association of social support networks, as well as the exacerbating association of work scheduling impact (WSI) with suicidal ideation, planning, and attempts of workers in the Australian entertainment industry. Findings indicated that mental health, WSI, and social support were significantly associated with workers’ suicide ideation in their lifetime (H_2). More importantly, high impact due to work scheduling (WSI) was shown to significantly associate with an increase in suicide ideation even in workers with good mental health. This highlights the disruptive association between working long and odd hours and suicidal behaviors among Australian

entertainment workers, denoting the need to regulate labor conditions in their industry appropriately [2, 7].

Alternatively, the moderating effect of WSI was not present in workers’ suicide ideation in their last 12 months (H_1), suicide planning (H_3), and suicide attempts (H_4). This denotes that (i) a more comprehensive approach to capture variables influencing suicide planning and attempts may be needed to increase our understanding of such behaviors, and (ii) that the exacerbating role of WSI on suicide ideation may not always be present during the initial 12 months of experiencing problems, due to working long and odd hours. In line with previous literature [26], this could imply that work-related stress may have a progressive or accumulated effect on one’s mental health and suicide ideation, which could become more tangible after 12 months of experiencing impaired sleep and disrupted social life associated to one’s work commitments.

Moderating effect of social support on the relationship between mental health and suicidal behaviors

Considering the buffering effect of social support on the relationship between mental health and suicidal behaviors, results provide mixed support to the proposed hypotheses. Specifically, support was only observed for our hypothesized association between lower social support and the relationship between mental health and suicide ideation in the last 12 months (H_5). This association is particularly important for workers of the Australian entertainment industry who may experience low social support

Table 2 Logistic regression coefficients with ‘Suicide planning’ and ‘Suicide attempts’ as outcome variables

Variable	B	BSE	Wald	p	Exp(B)	95% BCI for Exp(B)	
						Lower	Upper
Outcome Variable—Suicide planning							
α : Constant/Intercept	.735	1.840	.160	.689	2.086		
b_{31} : SF12	-.032	.065	.244	.622	.969	.853	1.099
b_{32} : WSI	-.013	.058	.047	.828	.987	.881	1.107
b_{33} : MSPSS	-.106	.307	.118	.731	.900	.493	1.643
b_{34} : SF12*WSI	.002	.003	.340	.560	1.002	.996	1.007
b_{35} : SF12*MSPSS	-.008	.013	.370	.543	.992	.967	1.018
b_{36} : SF12*WSI*MSPSS	.000	.001	.000	.999	1.000	.999	1.001
Outcome Variable—Suicide attempts							
α : Constant/Intercept	-.931	2.358	.156	.693	.394		
b_{41} : SF12	-.044	.086	.258	.611	.957	.808	1.133
b_{42} : WSI	.030	.074	.162	.688	1.030	.891	1.192
b_{43} : MSPSS	-.200	.390	.264	.608	.819	.381	1.757
b_{44} : SF12*WSI	.002	.004	.197	.657	1.002	.995	1.009
b_{45} : SF12*MSPSS	.004	.017	.063	.801	1.004	.971	1.038
b_{46} : SF12*WSI*MSPSS	.000	.001	.305	.581	1.000	.998	1.001

B Logistic coefficient, BSE Bootstrapped standard error, p Probability, Exp(B) Odds ratio, BCI Bootstrapped confidence interval, SF12 Short Form 12 items Mental Health Component, WSI Work Schedule Impact, MSPSS Multidimensional Scale of Perceived Social Support

considering their higher reported suicide ideation levels (Fig. 2 left panel). This is in line with past literature suggesting that increased levels of social support are associated with belongingness, social integration, feeling supported, and consequently improved mental wellbeing [2, 16, 19].

The buffering association of social support was not revealed considering suicide ideation lifetime (H_6), suicide planning (H_7), and suicide attempts (H_8). The lack of support for these hypotheses may indicate the potential importance of other individual and situational factors related to one's suicide behaviors beyond the level of social support experienced. Thus, it may be suggested that a 'systems approach' incorporating a range of protective and risk factors, at both the individual and the contextual level, as well as their interactions, should be considered to assess the transitioning from ideation to planning or execution and suicide behaviors in general [11]. Additionally, Klonsky and May [14] suggest that a progressive transition from moderate to severe suicide ideation usually occurs when hopelessness is greater than a sense of connectedness. This could explain the buffering association of social support on suicide ideation *only* during the first 12 months.

Moderated moderation – effect of WSI and social support on the relationship between mental health and suicidal behaviors

Finally, mixed results were observed considering the proposed interplay (moderated moderation effects) between WSI and social support regarding the relationship between mental health and suicidal behaviors among Australian entertainment industry workers. Specifically, our hypothesized role of the interaction between WSI and social support on the association between one's level of reported mental health and suicide ideation in the last 12 months was confirmed (H_9). Thus, one could suggest that while the nature of the entertainment industry may involve doing shift work (e.g., live performances are likely to happen at night or during the weekend), appropriately regulating the maximum number of working hours per day, providing fair remuneration (e.g., recognition of over-time) and compensation for working odd hours (e.g., time in lieu) may likely associate to weakening suicidal behaviors among more vulnerable workers. Indeed, past research identified *financial instability* as one of the main concerns associated with working in the Australian entertainment industry, signalling a power dynamic that may prevent workers from declining opportunities for paid work having to endure long and odd hours despite potentially adverse psychological consequences [2]. These recommendations seem particularly relevant considering that the effects of the Covid-19 pandemic may be largely unexplored, particularly in populations

whose financial security has been severely impacted due to lengthy periods of social restrictions.

Finally, it is important to denote the lack of support for proposed moderating effects associated with participants' suicide planning and attempts. This likely indicates that while both WSI and social networks are significantly associated with suicide behaviors, other personal traits/experiences are required for one to transition from suicidal ideation to planning and action. Indeed, Overholser and Spirito [27] propose that only a small proportion of individuals thinking about suicide attempt to take their life, with impulsivity and hopelessness due to the inability to improve situational contexts playing an important role. This highlights the importance of appropriately addressing differences in the assessment and prevention of suicide ideation, planning, and attempts [11]. Thus, additional cautiousness is recommended in the interpretation of the current findings.

Limitations and future research

Despite the value of these findings and the methodological strengths of the current study, limitations need to be considered. Firstly, given the cross-sectional nature of the employed data, collected in 2015, causative associations could only be speculated here, while potential changes during later developments may have not been accounted. Secondly, considering the inherent cultural elements of different countries, it is important to note that the findings presented in the current research might not be generalizable to workers of the entertainment industry in other parts of the world, and perhaps future research may seek to employ statistical methods to validate these results (i.e., measurement invariance [28, 29]). Thirdly, only self-report measures were used, increasing the risk of subjectivity biases. Hence, longitudinal studies assessing variables of relevance in workers of the entertainment industry of different countries while concurrently employing more objective measures (i.e., one's actual work schedule to assess WSI) would be recommended. Fourthly, considering that a time criterion was only employed to assess suicide ideation (i.e., 12 months vs. lifetime), a potential confounding effect of recent/lifetime suicide planning and attempts could play an important role in interpreting the associations explored here. Additionally, the question employed to address suicidal "planning" (i.e., "Have you planned to complete suicide?") may have been perceived by some participants as inquiring about one's intention to suicide without necessarily involving preparatory steps. Therefore, future studies may seek to clarify such potentially confounding results. Fifthly, the current study specifically

focused on the associations between psychological wellbeing, WSI and social support on suicidal behaviors; however, many other protective (e.g., coping strategies, or access to appropriate mental healthcare) and risk factors (e.g., income, alcohol and drugs, or previously diagnosed mental illness) influencing suicidal thoughts and behavior could be considered to provide a more systematic and more holistic approach to suicidal behaviors. For example, a prospective network analysis approach may enable simultaneously and comparatively assessing the effect of multiple such variables on suicidal behaviors [30]. Sixthly, the results reported here should be interpreted with caution, considering that suicide ideation was treated as a latent variable (i.e., composite score of three related questions) and limited empirical validity supports this methodological approach. Finally, the unexplored (in the present study) possibility of WSI operating as a moderator of the moderating influence of social support may also need to be explored in future research.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12888-022-04376-2>.

Additional file 1: Supplementary Table 1. Demographic statistics. **Supplementary Table 2.** Reliability Analysis. **Supplementary Table 3.** Exploratory Factor Analysis. **Supplementary Table 4.** Johnson and Neyman technique to determine the zone of significance.

Acknowledgements

We would like to acknowledge Dr Julie van den Eynde who led the project on working in the Australian Entertainment industry.

Authors' contributions

DZ contributed to the article's conceptualization, data curation, formal analysis, methodology, project administration, and writing of the original draft. CS contributed to the article's conceptualization, data curation, writing, review, and editing the final draft and project administration. AF contributed to data curation, writing, review, and editing the final draft and project administration. VS contributed to the review and edit of the final form of the manuscript. The author(s) read and approved the final manuscript.

Funding

The research was commissioned and funded by Entertainment Assist. VS has received the Australian Research Council, Discovery Early Career Researcher Grant/Award Number: DE210101107.

Availability of data and materials

The data that supports the findings of this study are available for Entertainment Assist, but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available for the authors upon reasonable request and with permission of Entertainment Assist. Please contact Daniel Zarate (daniel.zarate@live.vu.edu.au) to request access to the dataset.

Declarations

Ethics approval and consent to participate

Ethics approval granted by the Victoria University Ethics Committee. The current study only involved adult subjects (+18 years old) and informed consent

was obtained in all cases. All methods were carried out in accordance with relevant guidelines and regulations.

Consent for publication

Not applicable.

Competing interests

Dr Vasileios Stavropoulos is an associate editor for BMC Psychiatry. The remaining authors (Christopher Sonn, Adrian Fisher, and Daniel Zarate) declare no competing interests as defined by BMC, or other interests that might be perceived to influence the results and/or discussion in this paper.

Author details

¹Institute for Health and Sport, Victoria University, Melbourne, Australia. ²University of Athens, Athens, Greece.

Received: 26 June 2022 Accepted: 8 November 2022

Published online: 01 December 2022

References

- Kenny DT. Stairway to hell: Life and death in the pop music industry. *The Conversation*. 2014 <https://theconversation.com/stairway-to-hell-life-and-death-in-the-pop-music-industry-32735>
- Van den Eynde J, Fisher A, Sonn C. Working in the Australian Entertainment Industry: Final Report. Entertainment Assist, Victoria University. 2016 https://crewcare.org.au/images/downloads/WorkingintheAustralianEntertainmentIndCiteESMustry_FinalReport_Oct16.pdf
- Van Rens FECA, Heritage B. Mental health of circus artists: Psychological resilience, circus factors, and demographics predict depression, anxiety, stress, and flourishing. *Psychol Sport Exerc*. 2021;53:101850. <https://doi.org/10.1016/j.psychsport.2020.101850>.
- Gross SA, Musgrave G. Can Music Make you Sick? A Study into the Incidence of Musicians' Mental Health Part 2: Qualitative Study and Recommendations. University of Westminster / MusicTank. 2017 <https://www.helpmusicians.org.uk/assets/publications/files/can-music-make-you-sick-part-2-qualitative-study-1.pdf>
- Fujino Y, Horie S, Hoshuyama T, Tsutsui T, Tanaka Y. A systematic review of working hours and mental health burden. *J Occup Health*. 2006;48(4):87–97. <https://doi.org/10.1539/sangyoisei.48.87>.
- Flynn S, Nyathi T, Tham S-G, Williams A, Windfuhr K, Kapur N, Appleby L, Shaw J. Suicide by mental health in-patients under observation. *Psychol Med*. 2017;47(13):2238–45. <https://doi.org/10.1017/s0033291717000630>.
- King B, Berg L, Koenig J, Adair JJ, Tirado C. A Revised Occupational Stress Measure for Popular Musicians: Pilot Test of Validity and Reliability. *Med Probl Perform Art*. 2019;34(2):85–91. <https://doi.org/10.21091/mppa.2019.2015>.
- Preti A, De Basi F, Miotto P. Musical Creativity and Suicide. *Psychol Rep*. 2001;2001(89):719–27. <https://doi.org/10.2466/pr0.2001.89.3.719>.
- de Leo D, Goodfellow B, Silverman M, Berman A, Mann J, Arensman E, Hawton K, Phillips MR, Vijayakumar L, Andriessen K, Chavez-Hernandez A-M, Heisel M, Kölves K. International study of definitions of English-language terms for suicidal behaviours: a survey exploring preferred terminology. *BMJ Open*. 2021;11:e043409. <https://doi.org/10.1136/bmjopen-2020-043409>.
- Goodfellow B, Kölves K, de Leo D. Contemporary definitions of suicidal behavior: a systematic literature review. *Suicide Life Threat Behav*. 2019;49(2):488–504. <https://doi.org/10.1111/sltb.12457>.
- Klonsky ED, May AM, Saffer BY. Suicide, Suicide Attempts, and Suicidal Ideation. *Annu Rev Clin Psychol*. 2016;12(1):307–30. <https://doi.org/10.1146/annurev-clinpsy-021815-093204>.
- Crosby AE, Ortega L, Melanson C. Self-Directed Violence Surveillance: Uniform Definitions and Recommended Data Elements (Version 1.0). *CDC Natl Cent Inj Prev Control*; 2011. p. 1–96.
- Ribeiro JD, Franklin JC, Fox KR, Bentley KH, Kleiman EM, Chang BP, Nock MK. Self-injurious thoughts and behaviors as risk factors for future suicide ideation, attempts, and death: a meta-analysis of longitudinal studies. *Psychol Med*. 2015;46(2):225–36. <https://doi.org/10.1017/S003329171501804>.

14. Klonsky ED, May AM. The Three-Step Theory (3ST): a new theory of suicide rooted in the "ideation-to-action" framework. *Int J Cogn Ther*. 2015;8:114–29. <https://doi.org/10.1521/ijct.2015.8.2.114>.
15. Joiner, T. *Why People Die by Suicide*. Harvard University Press. 2005. <https://www.jstor.org/stable/j.ctvjghv2f>
16. Sonn CC, Fisher AT. Sense of Community: Community Resilient Responses to Oppression and Change. *J Community Psychol*. 1998;26(5):457–72. [https://doi.org/10.1002/\(SICI\)1520-6629\(199809\)26:5%3c457::AID-JCOP5%3e3.0.CO;2-O](https://doi.org/10.1002/(SICI)1520-6629(199809)26:5%3c457::AID-JCOP5%3e3.0.CO;2-O).
17. Pearson JE. The definition and measurement of social support. *J Couns Dev*. 1986;64(6):390–5. <https://doi.org/10.1002/j.1556-6676.1986.tb01144.x>.
18. Harandi TF, Taghinasab MM, Nayeri TD. The correlation of social support with mental health. A meta-analysis. *Electronic Physician*. 2017;9(9):5212–22. <https://doi.org/10.19082/5212>.
19. Huynh JY, Xanthopoulou D, Winefield AH. Social support moderates the impact of demands on burnout and organizational connectedness: A two-wave study of volunteer firefighters. *J Occup Health Psychol*. 2013;18(1):9–15. <https://doi.org/10.1037/a0030804>.
20. Crabtree JR. Tunesmiths and toxicity: workplace harassment in the contemporary music industries of Australia and New Zealand. *Open Publications of UTS Scholars*. 2020. <http://hdl.handle.net/10453/148011>
21. Faul F, Erdfelder E, Buchner A, Lang AG. Statistical power analyses using G* Power 3.1: Tests for correlation and regression analyses. *Behav Res methods*. 2009;41(4):1149–60.
22. Zimet GD, Dahlem NW, Zimet SG, Farley GK. The Multidimensional Scale of Perceived Social Support. *J Pers Assess*. 1988;52(1):30–41.
23. Revelle W. *Psych: Procedures for Psychological, Psychometric, and Personality Research*. [R package]. 2019. Retrieved from <https://cran.r-project.org/package=psych>
24. Ware JE, Kosinski M, Keller SD. *How to Score the SF-12 Physical and Mental Health Summary Scales* (2nd Ed.). The Health Institute, New England Medical Center. 1995. https://www.researchgate.net/publication/242636950_SF-12_How_to_Score_the_SF-12_Physical_and_Mental_Health_Summary_Scales
25. Hayes AF. *Introduction to mediation, moderation, and conditional process analysis: A regression-based approach*. Guildford Press; 2013.
26. Rosiek A, Rosiek-Kryszewska A, Leksowski L, Leksowski K. Chronic stress and suicidal thinking among medical students. *Environ Res Public Health*. 2016;13(2):212. <https://doi.org/10.3390/ijerph13020212>.
27. Overholser J, Spirito A. Precursors to adolescent suicide attempts. In: Spirito A, Overholser JC, Overholser J, editors. *Practical Resources for the Mental Health Professional: Evaluating and Treating Adolescent Suicide Attempters*. Academic Press; 2003. p. 19–40. <https://doi.org/10.1016/B978-012657951-2/50003-0>.
28. Zarate D, Marmara J, Potoczny C, Hosking W, Stavropoulos V. Body appreciation scale (BAS-2): measurement invariance across genders and item response theory examination. *BMC Psychol*. 2021;9(1):114. <https://doi.org/10.1186/s40359-021-00609-3>.
29. Marmara J, Zarate D, Vassallo J, Patten R, Stavropoulos V. Warwick Edinburgh mental well-being scale (WEMWBS): measurement invariance across genders and item response theory examination. *BMC Psychology*. 2022;10:31. <https://doi.org/10.1186/s40359-022-00720-z>.
30. Zarate D, Ball M, Montag C, Prokofieva M, Stavropoulos V. Unravelling the web of addictions: a network analysis approach. *Addict Behav Rep*. 2022;15(4):100406. <https://doi.org/10.1016/j.abrep.2022.100406>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

