

RESEARCH

Open Access



# Psychometric properties of the post-traumatic stress disorder checklist for DSM-5 (PCL-5) in Chinese stroke patients

Che Jiang<sup>1†</sup>, Gaici Xue<sup>1†</sup>, Shujing Yao<sup>1</sup>, Xiwu Zhang<sup>1</sup>, Wei Chen<sup>2</sup>, Kuihong Cheng<sup>1</sup>, Yibo Zhang<sup>1</sup>, Zhensheng Li<sup>3</sup>, Gang Zhao<sup>1</sup>, Xifu Zheng<sup>2</sup> and Hongmin Bai<sup>1\*</sup>

## Abstract

**Background** Stroke is a devastating disease and can be sufficiently traumatic to induce post-traumatic stress disorder (PTSD). Post-stroke PTSD is attracting increasing attention, but there was no study assessing the psychometric properties of the PCL-5 in stroke populations. Our study was conducted to examine the psychometric properties of the PTSD Checklist for DSM-5 (PCL-5) in Chinese stroke patients.

**Methods** This was a cross-sectional observational study conducted at our hospital. Three hundred and forty-eight Chinese stroke patients came to our hospital for outpatient service were recruited. They were instructed to complete the PCL-5 scales and were interviewed for PTSD diagnosis with the Clinician Administered PTSD Scale for DSM-5 (CAPS-5). The cutoff scores, reliability and validity of the PCL-5 were analyzed.

**Results** PCL-5 scores in our sample were positively skewed, suggesting low levels of PTSD symptoms. The reliability of PCL-5 was good. Exploratory and confirmatory factor analyses indicated acceptable construct validity, and confirmed the multi-dimensionality of the PCL-5. By CFA analysis, the seven-factor hybrid model demonstrated the best model fit. The PCL-5 also showed good convergent validity and discriminant validity. Receiver operating characteristic (ROC) analyses revealed a PCL-5 score of 37 achieved optimal sensitivity and specificity for detecting PTSD.

**Conclusions** Our findings supported the use of PCL-5 as a psychometrically adequate measure of post-stroke PTSD in the Chinese patients.

**Keywords** Stroke, PTSD, PCL-5, CAPS-5, Psychometric property

## Background

Post-traumatic stress disorder (PTSD) is a psychiatric disorder which develops as a result of exposure to life-threatening traumatic events. According to the Statistical Manual of Mental Disorders-version 5 (DSM-5), PTSD is characterized by four clusters of symptoms: recurrent involuntary intrusive memories, avoidance, negative alterations in cognition and mood, and alterations in arousal and reactivity [1]. The PTSD Checklist (PCL) is one of the most frequently used self-report screener for PTSD, and the PCL-5 is the updated version that conforms to the DSM-5 criteria [2]. Compared to PCL-M/PCL-C/

<sup>†</sup>Che Jiang and Gaici Xue contributed equally to this work and share first authorship.

\*Correspondence:

Hongmin Bai  
baihmfmmu@vip.163.com

<sup>1</sup> Department of Neurosurgery, General Hospital of Southern Theatre Command, 111 Liuhua Road, Guangzhou 510010, China

<sup>2</sup> School of Psychology, South China Normal University, Guangzhou 510631, China

<sup>3</sup> Department of Neurology, General Hospital of Southern Theatre Command, Guangzhou 510010, China



PCL-S which conforms to the DSM-IV criteria, PCL-5 is the only version designed to evaluate the severity of PTSD symptoms across various types of traumas. It has been translated to several languages and showed good internal consistency, test–retest reliability, and validity in different samples [3–6]. The Chinese version of PCL-5 has been translated and used in traumatized samples [7–9].

Most previous studies focused on the PTSD secondary to external traumatic events such as combat and sexual violence, while less studies focused on medical conditions or acute illness [4, 10, 11]. Stroke is a serious health event and presents one of the leading causes of death and disability in the worldwide [12]. It has subtypes like cerebral infarction, intracerebral hemorrhage, transient ischemic attack (TIA), and subarachnoid hemorrhage (SAH). According to a meta-analysis on post-stroke mood disorder, the prevalence of major depressive disorder, anxiety, dysthymia, and adjustment disorder were 17.7%, 9.8%, 3.1%, and 6.9% respectively [13]. Stroke features sudden onset of neurologic deficits and potentially threatens patients' lives, which kind of medical events is qualified as traumatic events according to the DSM-5 criteria. Due to sample heterogeneity, the prevalence of PTSD (or PTSD symptom) varies greatly, ranging from 4 to 37% [14]. Although some studies showed that the degree of disability caused by stroke correlates positively with PTSD symptoms [15–18], mild stroke [19] and TIA [20] can also lead to PTSD.

China bears the biggest stroke burden in the world, and stroke is associated with the highest disability-adjusted life-years lost in this country [21]. Annual estimates of 11 million prevalent cases and 2.4 million new cases were reported [22]. Due to the recurrent nature of stroke, long-term medical management is especially important. The comorbidity of PTSD is related to stroke patients' non-adherence of treatment [23, 24], as well as lower quality of life [25, 26]. So, regular psychological assessments using PCL-5 at follow-ups for stroke patients are helpful and deserves wider application. Presently, there are no studies focusing on the psychometric properties of PCL-5 in stroke patients, thus our study aimed at solving this problem in the Chinese sample. Specifically, the evaluation goals were to explore the reliability of the PCL-5, the factor loadings and construct validity by factor analyses, and the convergent and discriminant validity through correlations between the PCL-5 and another measure of PTSD (i.e., CAPS-5) and between the PCL-5 and measures of other constructs (e.g., anxiety, depression, and alcohol abuse), respectively.

## Methods

### Participants

All methods were carried out in accordance with relevant guidelines and regulations. The study was approved by the Ethics Committee of General Hospital of Southern Theatre Command. Chinese stroke patients came to our hospital (PLA General Hospital of Southern Theatre Command) for outpatient service from August 2021 to August 2022 were asked to participate in this survey if the doctors regarded the patients can cooperate. The exclusion criteria were as follows: age < 18 years or > 80 years; severe aphasia or cognitive impairment; refused to participate in this survey. There were no financial or other incentives for taking part, except for the doctor's encouragement. Over 700 patients were initially invited. Finally, three hundred and forty-eight patients were included, but the rest either refused due to various reasons or were ineligible. All of the participants gave their written consent for participation. The mean age of our sample was 55.41, older than that of most samples in other researches on PTSD. The majority of our participants were male, with a high school or lower educational level, and not in employment. Cerebral infarction represented the major subtype of stroke (Table 1).

### Procedure

All participants were asked to complete self-reported scales, and their demographic data were obtained. Further explanation of any items in the PCL-5 were not given by specialists even if participants asked. Then, participants were interviewed using CAPS-5 by masters- and doctoral-level clinicians with formal training on the measure. The assigned interviewers took part in CAPS-5 training workshop and were supervised throughout the study. The training and supervision were conducted by the psychologists listed as the fifth and the second to last authors. The participants were instructed to rate their symptoms only in relation to their stroke experience but not to other traumatic events when completing the PCL-5 and CAPS-5. The CAPS-5 was administered by interviewers without knowledge of the PCL-5 scores. Twenty interviews were audio-recorded for the evaluation of inter-rater reliability. After the survey, participants were thanked for their participation.

### Measures

The Chinese versions of all measures used in our study were adapted by a two-stage process of translation and back translation by bilingual researchers based on the original English versions, and they were further verified by senior experts.

**Table 1** Demographic data

	<i>M</i>	<i>SD</i>
Age	55.41	10.58
NIHSS score	2.69	2.62
	<i>Count</i>	<i>Percent</i>
Gender (female)	94	27.01%
Marital status		
Married or cohabiting	255	73.28%
Single	85	24.43%
Educational level		
High school or lower	278	79.89%
University or higher	65	18.68%
Employment status		
Employed	126	36.21%
Unemployed/retired	220	63.22%
Stroke type		
CI	269	77.30%
IH	33	9.48%
TIA	39	11.21%
Other	7	2.01%
Lesion site		
Cerebrum	209	60.06%
Cerebellum	22	6.32%
Brainstem	39	11.21%
Other/unknown	78	22.41%
Course of stroke		
< 0.5 year	69	19.83%
0.5–1 year	95	27.30%
> 1 year	184	52.87%

Values are expressed as mean ± SD or percentage

CI Cerebral infarction, IH Intracerebral hemorrhage, TIA Transient ischemic attack, SAH Subarachnoid hemorrhage

**Hospital Anxiety and Depression scale (HADS) [27]**

This scale has two subscales of depression (HADS-D) and anxiety (HADS-A), each containing 7 items. The items are rated from 0 (most of the time) to 3 (absolutely not). Higher scores indicate greater degree of anxiety or depression. A subscale score of ≥ 8 suggests being anxious or depressed, respectively. The Chinese version showed good internal and test–retest reliability [28–30].

**Alcohol Use Disorder Identification Test (AUDIT)**

This scale is used to assess the habits of alcohol consumption and related hazardousness [31]. It has 10 items and its total score ranges from 0 to 40. Higher score indicates worse alcohol related problems. A total score of ≥ 8 indicates hazardous or harmful drinking. Good psychometric properties have been proved for the Chinese translated version [5, 6, 32–34].

**Post-traumatic stress disorder checklist for DSM-5 (PCL-5)**

The PCL-5 is a 20-item self-report scale evaluating DSM-5 PTSD symptoms [35]. Items 1–5 on the PCL-5 correspond to Criterion B, items 6–7 to Criterion C, items 8–14 to Criterion D, and items 15–20 to Criterion E. The 20 item is rated on a 5-point Likert scale (0 = not at all, 1 = a little bit, 2 = moderately, 3 = quite a bit and 4 = extremely) reflecting the severity of traumatic experience. The scores are summed to create a total severity score (range 0–80). This scale has been validated or use in cardiovascular populations [36] and been frequently used in assessing stroke patients [37, 38]. The Chinese version has been validated and widely used [39–41].

**The clinician-administered PTSD scale for DSM-5 (CAPS-5) [42]**

The CAPS-5 is a scale designed for the diagnosis of PTSD, and is used for semi-structured interview. Its items correspond to DSM-5 criteria of PTSD (criterion A to H), and includes 5 items for criterion B, 2 for criterion C, 7 for criterion D, and 6 for criterion E. Each of the items (criterion B to E) is rated on a Likert (0–4) scale from 0 = absent to 4 = extremely, with a total score ranging from 0 to 80. According to the SEV2 rule for CAPS-5 symptom scoring, a severity rating of 2 (moderate) or higher indicates presence of a symptom or impairment [43]. A PTSD diagnosis is made based on the participant satisfying all DSM-5 criteria, fulfilling at least one item each of intrusion (criterion B) and avoidance (criterion C), two items each of negative changes in cognitions and mood (criterion D) and hyperarousal (criterion E), functional impairment (criterion G) and the presence of symptoms for ≥ 1 month (criterion F). Psychometric evaluation of the CAPS-5 demonstrated good internal consistency, strong inter-rater reliability and test–retest reliability [43].

**Statistical analysis**

Data were analyzed using SPSS 18.0 and AMOS 22.0. The skewness for PCL-5 items were first psychometrically examined. The reliability of PCL-5 and other measures were measured. Furthermore, item-total correlations and Cronbach’s alpha-if item deleted for the PCL-5 were reported. Exploratory factor analysis (EFA) was conducted to examine the factor loadings and construct validity (the factors that have values greater than 1 are extracted according to Kaiser-Guttman criterion), and confirmatory factor analysis (CFA) was performed to determine the construct validity of the four-factor DSM-5 model (intrusion, avoidance, negative alterations in cognitions and mood, and hyperarousal), the six-factor anhedonia model (Intrusion, avoidance,

negative affect, anhedonia, dysphoric arousal, anxious arousal) [41], and the seven-factor hybrid model of PTSD (Intrusion, avoidance, negative affect, anhedonia, externalizing behaviors, anxiety arousal, dysphoric arousal) [44]. The EFA analysis was conducted using principal component analysis (PCA), with varimax axis rotation and Kaiser normalization. The factor loading of each item in EFA is regarded as acceptable if >0.5. By CFA, the chi-square, df, chi-square/df, *p*, goodness of fit index (GFI), CFI (comparative fit index), TLI (Tucker-Lewis index), AIC (Akaike information criterion), RMSEA (mean square error of approximation), and SRMR (standardized root mean square residual) were evaluated. Typically, GFI > 0.8 [45], RMSEA < 0.1 and SRMR < 0.08 [46] indicate acceptable model fit, whereas CFI > 0.9 and TLI > 0.9 indicate adequate model fit [47]. Convergent validity was evaluated with Spearman correlations between scores of PCL-5 and CAPS-5, and discriminant validity was tested with Spearman correlations between scores of PCL-5 and questionnaires of anxiety (HADS-A), depression (HADS-D), and alcohol abuse (AUDIT). The diagnostic utility of PCL-5 was calculated with Receiver Operating Curve (ROC) analysis. Specifically, Positive Predictive Power (PPP), Negative Predictive Power (NPP), Overall Correct Classification (OCC),  $\kappa$  (0.5) were reported.

Kappa statistics were considered fair agreement for 0.21–0.40, moderate for 0.41 to 0.60, substantial for 0.61 to 0.80, and almost perfect for 0.81–1.0 [48]. The highest value of sensitivity plus specificity was used to determine the optimal cut-off score.

**Results**

Descriptive analyses for PCL-5 items indicated positive skewness and leptokurtic (Table 2). The item-total correlation contained no negative values, indicating that the items were assessing the same construct. The descriptive characteristics and internal consistency of PCL-5 and other measures were shown in Table 3. The probable prevalence of anxiety (HADS-A score ≥ 8), depression (HADS-D score ≥ 8) and problem alcohol use (AUDIT score ≥ 8) after stroke attack were 18.97%, 12.93% and 8.91%, respectively. Fourteen participants (4.02%) were formally diagnosed with PTSD through interview. In our stroke patient sample, the overall severity of PTSD symptoms was low and the data were positively skewed. The total scores and all four factors showed good internal consistency (all Cronbach’s  $\alpha$  > 0.8 and Guttman’s split-half > 0.7).

After checked for audibility and completeness of the 20 audio-recorded CAPS-5 interviews, 19 appeared to be appropriate for the second rating. The inter-rater

**Table 2** Item-level psychometric properties of the PCL-5

PCL-5 Item	M	SD	Variance	Skew	Kurtosis	Range	Item-total correlation	Cronbach’s $\alpha$ if item deleted
1 (B1)	0.68	1.11	1.23	1.60	1.54	0–4	0.72	0.93
2 (B2)	0.68	1.04	1.08	1.74	2.47	0–4	0.61	0.93
3 (B3)	0.60	0.90	0.82	1.69	2.63	0–4	0.62	0.93
4 (B4)	0.78	0.97	0.93	1.34	1.54	0–4	0.66	0.93
5 (B5)	0.37	0.77	0.59	2.60	7.53	0–4	0.50	0.93
6 (C1)	0.66	1.05	1.10	1.69	2.13	0–4	0.56	0.93
7 (C2)	0.60	1.00	0.99	1.76	2.49	0–4	0.64	0.93
8 (D1)	0.29	0.62	0.38	2.71	9.25	0–4	0.45	0.93
9 (D2)	0.42	0.76	0.58	2.30	5.92	0–4	0.53	0.93
10 (D3)	0.51	0.82	0.68	2.02	4.58	0–4	0.51	0.93
11 (D4)	0.77	0.99	0.99	1.49	2.07	0–4	0.44	0.93
12 (D5)	0.75	1.02	1.04	1.50	1.85	0–4	0.60	0.93
13 (D6)	0.56	0.88	0.77	1.79	3.19	0–4	0.62	0.93
14 (D7)	0.53	0.84	0.71	2.17	5.58	0–4	0.52	0.93
15 (E1)	0.53	0.73	0.54	1.50	2.66	0–4	0.56	0.93
16 (E2)	0.32	0.65	0.43	2.41	6.62	0–4	0.51	0.93
17 (E3)	0.57	0.78	0.60	1.88	4.90	0–4	0.49	0.93
18 (E4)	0.43	0.77	0.59	2.23	5.69	0–4	0.53	0.93
19 (E5)	0.56	0.82	0.67	2.06	5.35	0–4	0.46	0.93
20 (E6)	0.93	1.09	1.19	1.34	1.30	0–4	0.50	0.93

**Table 3** Descriptive characteristics and internal consistency of PCL-5 and other measures

	M	SD	Cronbach's $\alpha$	Guttman's split-half	N
Intrusion	3.11	3.90	0.87	0.78	5
Avoidance	1.27	1.88	0.82	0.82	2
Negative alterations in cognition and mood	3.83	4.07	0.81	0.72	7
Alterations in arousal and reactivity	3.35	3.58	0.83	0.82	6
Total PCL-5 score	11.56	11.90	0.94	0.91	20
CAPS-5 score	8.92	10.11	0.92	0.91	20
HADS-A score	4.87	3.82	0.79	0.78	7
HADS-D score	3.99	3.12	0.77	0.77	7
AUDIT	2.54	3.21	0.79	0.84	10

reliability was high for total PTSD symptom severity score ( $ICC=0.97$ ,  $95\% CI=0.92-0.99$ ), and was perfect for PTSD diagnosis ( $\kappa=1.0$ ) with agreement on all classifications between raters.

We performed the EFA to see the factor loadings of each item and whether the theoretically expected four factors can be observed in our sample. This sample was suitable for EFA analysis (Kaiser Meyer Olkin = 0.95, chi-square = 3531.97,  $df=190$ ,  $p < 0.001$ ). Results showed that the four new factors explained 62.02% of total variance indicating a good representation of the whole data. The factor loadings were presented in Table 4. The analysis demonstrated four factors, although slightly different from the theoretical factor structure. The one to four factors explained 20.68%, 15.71%, 12.96%, and 12.67% of the total variance, respectively. The first factor primarily represented intrusion (4 items) plus avoidance factor (2 items). The second factor primarily represented the factor of alterations in arousal and reactivity (6 items). The factor of negative alterations in cognitions and mood was basically divided into the third (3 items) and fourth factors (3 items). The factor loadings of most items were good (factor loading > 0.5). After removing the 3 items with factor loadings < 0.5 (loss of interest, hypervigilance, and concentration), the factors can explain 65.04% of total variance, slightly elevated compared to the original 20 items.

CFA were then used to evaluate the PCL-5 with original four factors (Fig. 1; Table 5). Factor loadings of all item were acceptable (ranged from 0.55 to


**Table 4** Factor loadings of PCL-5 items

PCL-5 Item	Item Description	Factors			
		1	2	3	4
1(B1)	Memories	<b>0.707</b>	0.187	0.449	0.219
2(B2)	Dreams	<b>0.794</b>	0.212	0.181	0.128
3(B3)	Flashbacks	<b>0.699</b>	0.082	0.305	0.225
4(B4)	Cued distress	0.451	0.172	<b>0.617</b>	0.189
5(B5)	Cued physical reactions	<b>0.689</b>	0.248	0.160	0.168
6(C1)	Avoiding memories	<b>0.624</b>	0.377	0.170	0.281
7(C2)	Avoiding external reminders	<b>0.639</b>	0.335	0.128	0.314
8(D1)	Dissociative amnesia	0.120	<b>0.523</b>	0.309	0.246
9(D2)	Negative beliefs	0.217	0.289	0.162	<b>0.678</b>
10(D3)	Blame	0.153	0.177	0.152	<b>0.770</b>
11(D4)	Negative feelings	0.239	0.118	0.111	<b>0.724</b>
12(D5)	Loss of interest	0.373	0.212	<b>0.426</b>	0.396
13(D6)	Detachment/estrangement	0.263	0.297	<b>0.703</b>	0.116
14(D7)	Numbing	0.172	0.158	<b>0.838</b>	0.135
15(E1)	Irritability or aggressive behavior	0.137	<b>0.773</b>	0.135	0.109
16(E2)	Reckless behavior	0.145	<b>0.614</b>	0.409	0.269
17(E3)	Hypervigilance	0.420	<b>0.465</b>	0.232	0.176
18(E4)	Exaggerated startle	0.448	<b>0.658</b>	0.105	0.057
19(E5)	Concentration	0.387	<b>0.444</b>	0.139	0.356
20(E6)	Sleep	0.312	<b>0.572</b>	0.092	0.326

0.88). Parameters such as chi-square, GFI (> 0.8), CFI (> 0.9), TLI (> 0.9), AIC, RMSEA (< 0.08), standardized RMR (= 0.048) indicated good model fit [45–47, 49]. The six-factor anhedonia model and the seven-factor hybrid model of PTSD were also assessed by CFA and both showed better model fit than the original four-factor model, with the hybrid model demonstrated the best (Figs. 2 and 3; Table 5).

By Spearman correlation test, PCL-5 total scores and subscale scores all showed moderate to strong positive correlations with PTSD criteria in CAPS-5 (Table 6). This implied good convergent validity of PCL-5. PCL-5 total score also showed strong positive correlations with total scores of HADS-A and HADS-D for the entire sample, however, it did not demonstrate correlations with AUDIT (Table 7).

By ROC analysis, the PCL-5 total score's area under the curve (AUC) was 0.99 (SE = 0.003) indicating a very good level of diagnostic accuracy. The optimal cutoff scores of PCL-5 for the PTSD diagnosis as assessed through the CAPS-5 in our sample was 37, which had the highest sensitivity and specificity (Table 8).



**Fig. 1** Standardized estimation in CFA analysis results of PCL-5 of the four-factor DSM-5 model. Note. GFI, goodness of fit index; TLI, TuckerLewis index-in comparison with the null model; AIC, Akaike information criterion-in comparison with the null model; CFI, comparative fix index; RMSEA, root mean square error of approximation

**Discussion**

Although many scales have been designed for assessing PTSD symptoms, there lacks a reliable and valid screening instrument specifically designed for post-stroke PTSD, which has been increasingly drawing interest. Our study was the first to evaluate the psychometric properties of the PCL-5 among the

Chinese stroke patients. Our results indicated the PCL-5 has a good factor structure in assessing the current sample. Additionally, we found that the overall PTSD symptoms in our sample was relatively low with a 4% of diagnosed PTSD rate. A cutoff score of 37 for diagnostic value of PCL-5 was suggested by our study.

**Table 5** Confirmatory factor analysis of different models of PCL-5

	DSM-5 model	Anhedonia model	Hybrid model
Chi-square ( $\chi^2$ )	428.238	364.026	333.800
df	164	155	149
Chi-square/df	2.611	2.349	2.240
$p$	< 0.001	< 0.001	< 0.001
Goodness of Fit Index (GFI)	0.886	0.904	0.911
Comparative Fit Index (CFI)	0.923	0.939	0.946
Tucker-Lewis Index (TLI)	0.910	0.925	0.931
Akaike Information Criterion (AIC)	520.238	474.026	455.800
Mean Square Error of Approximation (RMSEA)	0.068	0.062	0.060
Standardized Root Mean Square Residual (SRMR)	0.0478	0.0454	0.0433


According to previous studies, prevalence or incidence of post-stroke PTSD varied greatly (from 4 to 37%) [14]. The PTSD rate and symptom severity in our sample were relatively low. This may be due to the different social and cultural background of the Chinese population. Moreover, participants in our sample were all outpatients, potentially excluding those with severe psychiatric or neurologic symptoms. Also, the majority of participants were male (73%), who may be less prone to develop PTSD [25, 50]. The selection bias may lead to milder PTSD symptoms in our study. Additionally, despite heterogeneity in methodology, previous small sample studies indicated the PTSD symptoms after stroke gradually decreased with the times. A meta-analysis in revealed the estimated rate of PTSD/PTSS following stroke or TIA was 23% within 1 year and 11% after 1 year [51]. More than half of our participants had stroke courses longer than 1 year. Future studies with large sample size and longitudinal design are required to further elucidate the temporal variation of PTSD rates after stroke and the possibility of delayed-onset PTSD.

In previous studies of other populations, the Cronbach's alpha value of PCL-5 total score ranged from 0.91 to 0.97, with that of individual item ranged from 0.79 to 0.93 [4–6]. The Cronbach' alpha in our sample was also excellent (total score, 0.94; individual item, 0.81 to 0.87).

Since models of EFA and CFA are different, previous studies showed that factor structures obtained by EFA often turn out to fit poorly in CFA [4, 52]. Thus we conducted both analyses. In the EFA, four factor model was detected, but was different from the theoretical model. The most prominent structure discrepancies were that the factor of avoidance was integrated into the factor of

intrusion, and the factor of negative alterations in cognitions and mood was split into two factors. This result was quite similar to the previous study on Greek women after cesarean section [4]. The possibility of symptom overlaps or interdependence of the two symptom clusters of intrusion and avoidance has been proposed by some previous studies in different samples [6, 53]. In a study of Chinese adolescents surviving an earthquake, analyses similarly split the PCL-5 factor of negative alterations in cognitions and mood into negative affect factor and anhedonia factor [54]. Our results were consistent with this study. In our sample, after deleting the factors with loadings < 0.5, the total variance explained only increased slightly (by 3.02%), thus the factor's structure was reserved.

According to EFA, loss of interest ("Loss of interest in activities that you used to enjoy?"), hypervigilance ("Being "superalert" or watchful or on guard?") and concentration ("Having difficulty concentrating?") are items with factor loadings of < 0.5. This may be explained by the specificity of stroke. Stroke often causes neurologic deficits such as limb weakness, bradylalia, and dizziness, which may very well lead to loss of some interests (e.g., sports, drinking, and socializing); Stroke has the potential of recurrence and exaggeration, and it may be effectively treated with thrombolysis or thrombectomy if detected in the super early stage after attack. Thus, patients are generally taught to be alert on the initial evidence of recurrence; Additionally, stroke patients are commonly individuals with older ages (averaging 55.41 years old in our sample) who have reduced energy compared to younger ones, and the lesion in brain may more or less influence cognition and concentration. These above symptoms are the results of stroke itself and may not represent PTSD symptoms.




**Fig. 2** Standardized estimation in CFA analysis results of PCL-5 of the six-factor anhedonia model. Note. GFI, goodness of fit index; TLI, TuckerLewis index-in comparison with the null model; AIC, Akaike information criterion-in comparison with the null model; CFI, comparative fix index; RMSEA, root mean square error of approximation

Through CFA analysis, the PCL-5 also conformed to the original dimensions, meaning the four-factor structure worked adequately for the Chinese stroke patients. Besides the DSM-5 model, previous studies also proposed other models such as Anhedonia model [41] and hybrid model [44] for PTSD. In our study, the seven-factor hybrid model exhibited the best fit. This result was in accordance with the findings on diverse populations [6, 40, 55–60]. Thus, whether the DSM-5 model remains

the best fit in specific populations is left for future research. In our sample, the PCL-5 scores demonstrated good convergent validity with CAPS-5 criteria, with correlations ranging from 0.568 to 0.747. Additionally, the PCL-5 scores also showed good discriminant validity by being strongly correlated with scales evaluating anxiety (HADS-A) and depression (HADS-D) while being not correlated with alcohol habits (AUDIT). These results were consistent with previous studies, except for





**Fig. 3** Standardized estimation in CFA analysis results of PCL-5 of the seven-factor hybrid model. Note. GFI, goodness of fit index; TLI, TuckerLewis index-in comparison with the null model; AIC, Akaike information criterion-in comparison with the null model; CFI, comparative fix index; RMSEA, root mean square error of approximation

**Table 6** Correlations among PCL-5 factors and CAPS-5 criterions (Bivariate Spearman rho)

	Intrusion	Avoidance	Negative alterations in cognitions and mood	Alterations in arousal and reactivity	PCL-5 total score
CAPS-5 B	.742**	.549**	.546**	.483**	.711**
CAPS-5 C	.579**	.747**	.487**	.418**	.632**
CAPS-5 D	.520**	.449**	.633**	.453**	.600**
CAPS-5 E	.495**	.466**	.439**	.568**	.539**

\*\* P < 0.01 (2-tailed)

**Table 7** Discriminant validity correlations for the PCL-5 and other measures

Measure	1	2	3	4	5
1. PCL-5	1.00				
2. CAPS-5	0.85**	1.00			
3. HADS-A	0.73**	0.68**	1.00		
4. HADS-D	0.74**	0.66**	0.58**	1.00	
5. AUDIT	0.07	0.07	0.09	0.04	1.00

\*\*  $P < 0.01$  (2-tailed)

the weak correlation between PCL-5 and AUDIT [5, 6]. This may be due to the reason that stroke patients generally quit drinking as advised by doctors and most of the participants in our sample only had mild or no PTSD symptoms, not severe enough to influence drinking habits.

For the Chinese version of the PCL-5, a cutoff point of 37 was the one that had the best predictive values (Youden index  $J = 0.979$ ). According to previous studies, the cutoff scores of PCL-5 was suggested ranging from 25 to 45 [5, 6, 61–64] in different populations. Our indicated value of 37 was average compared to these studies. Since mental disorders may negatively affect stroke patients’ prognosis, it is critical to screen PTSD in this population. However, due to the large number of stroke patients, there lacks enough neurological psychologists/psychiatrists for PTSD interview. Our study suggested that the self-report PCL-5 can fulfill the need of initial screening. In this sample a cut-off of 37 provided a sensitivity of 1.00 and

a specificity of 0.979, demonstrating a high degree of diagnostic accuracy. It should be pointed out that whether and at what level to set a cut-off and caseness scores is based on the purpose for using the scale [65]. It is also of important clinical value to detect individuals reporting significant subthreshold symptoms, since there are instances where treatment may benefit these individuals who are experiencing clinically significant functional impairment [6]. With the popularization and application of PCL-5 for initial screening and follow-up assessment, PTSD symptoms can be easily monitored facilitating future studies on the early intervention for post-stroke PTSD.

There were some limitations of our study. First, in our sample, the proportion of formally diagnosed PTSD was small and the overall PTSD symptoms was low. This may influence the analysis of cutoff value. Studies on samples with higher rate of PTSD patients are required in the future. Second, since social and cultural background can influence the psychometric properties, our results may not necessarily be translated to stroke populations in other races or societies. Third, because previous studies showed that PCL-5 is correlated with alcohol abuse [66, 67], we initially included AUDIT. However, the use of the AUDIT scale has not been validated in stroke patients, thus its application as a criterion for convergent and divergent validity may be debated. Fourth, in the study, we only aimed to investigate the psychometric properties of measures associated with PTSD, so we did not collect information on history of psychiatric disorders, which may be an important exclusion criterion in studies on epidemiological

**Table 8** Various PCL-5 cutoff scores in predicting PTSD diagnosis with DSM-5 standard

Cutoff score	Sensitivity	Specificity	PPP	NPP	OCC	$\kappa$ (0.5)
31.5	1.000	.949	0.438	1	0.948	0.585
32.5	1.000	.955	0.483	1	0.957	0.631
33.5	1.000	.970	0.583	1	0.971	0.723
34.5	1.000	.973	0.609	1	0.974	0.744
35.5	1.000	.976	0.636	1	0.977	0.766
37.0	1.000	.979	0.667	1	0.980	0.790
38.5	.929	.982	0.684	0.997	0.980	0.778
40.5	.929	.985	0.722	0.997	0.983	0.804
42.5	.929	.988	0.765	0.997	0.986	0.831
43.5	.857	.991	0.80	0.994	0.986	0.820
44.5	.714	.991	0.769	0.988	0.980	0.730
45.5	.714	.994	0.833	0.988	0.983	0.760

PPP (Positive Predictive Power) = true positives / (true positives + false positives); NPP (Negative Predictive Power) = true negatives / (true negatives + false negatives); OCC (Overall Correct Classification) = (true positives + true negatives) / (true positives + true negatives + false positives + false negatives);  $\kappa$  (0.5) = quality of efficiency

studies on post-stroke PTSD, revealing its prevalence and risk factors.

## Conclusion

Our findings support use of the PCL-5 as a psychometrically adequate measure of post-stroke PTSD in the Chinese patients.

## Abbreviations

PTSD	Post-traumatic stress disorder
PCL-5	Post-traumatic Stress Disorder (PTSD) Checklist for DSM-5
CAPS-5	Clinician Administered PTSD Scale for DSM-5
ROC	Receiver operating characteristic
TIA	Transient ischemic attack
SAH	Subarachnoid hemorrhage
EFA	Exploratory factor analysis
CFA	Confirmatory factor analysis

## Acknowledgements

None

## Authors' contributions

HB and CJ designed the study. CJ and GX wrote the manuscript. WC select proper questionnaires. SY, KC, and YZ conducted the survey. ZL analyzed the data. GZ and Xiwu Zhang extracted the data. Xifu Zheng revised the manuscript. The author(s) read and approved the final manuscript.

## Funding

This work was supported by the Military Medical Science and Technology Youth Cultivate Program under Grant [number 20QNPY078] and the Science and Technology Planning Project of Guangzhou under Grant [number 202102021268].

## Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author [HB] on reasonable request.

## Declarations

### Ethics approval and consent to participate

All methods were carried out in accordance with relevant guidelines and regulations. The study was approved by the Ethics Committee of General Hospital of Southern Theatre Command (No. 2020–55). Informed consent was obtained from all subjects.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

Received: 29 September 2022 Accepted: 23 December 2022

Published online: 09 January 2023

## References

- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. Arlington, VA: Author; 2013.
- Weathers FW, Litz BT, Keane TM, Palmieri PA, Marx BP, Schnurr PP. (2013). The PTSD checklist for DSM-5 (PCL-5). Scale available from the National Center for PTSD at [www.ptsd.va.gov](http://www.ptsd.va.gov).
- Bockhop F, Zeldovich M, Cunitz K, et al. Measurement invariance of six language versions of the post-traumatic stress disorder checklist for DSM-5 in civilians after traumatic brain injury. *Sci Rep*. 2022;12(1):16571.
- Orovou E, Theodoropoulou IM, Antoniou E. Psychometric properties of the post traumatic stress disorder checklist for DSM-5 (PCL-5) in Greek women after cesarean section. *PLoS ONE*. 2021;16(8): e0255689.
- Roberts NP, Kitchiner NJ, Lewis CE, Downes AJ, Bisson JI. Psychometric properties of the PTSD checklist for DSM-5 in a sample of trauma exposed mental health service users. *Eur J Psychotraumatol*. 2021;12(1):1863578.
- Morrison K, Su S, Keck M, Beidel DC. Psychometric properties of the PCL-5 in a sample of first responders. *J Anxiety Disord*. 2021;77: 102339.
- Li G, Wang L, Cao C, et al. DSM-5 posttraumatic stress symptom dimensions and health-related quality of life among Chinese earthquake survivors. *Eur J Psychotraumatol*. 2018;9(1):1468710.
- Jiang C. Risk of probable posttraumatic stress disorder in patients with trigeminal neuralgia. *Psychol Health Med*. 2019;24(4):493–504.
- Wang L, Cao X, Cao C, Fang R, Yang H, Elhai JD. Factor structure of DSM-5 PTSD symptoms in trauma-exposed adolescents: Examining stability across time. *J Anxiety Disord*. 2017;52:88–94.
- Jiang C, Wang J. Post-traumatic stress disorders in patients with low-grade glioma and its association with survival. *J Neurooncol*. 2019;142(2):385–92.
- Edmondson D, von Känel R. Post-traumatic stress disorder and cardiovascular disease. *Lancet Psychiatry*. 2017;4(4):320–9.
- Béjot Y, Daubail B, Giroud M. Epidemiology of stroke and transient ischemic attacks: current knowledge and perspectives. *Rev Neurol (Paris)*. 2016;172(1):59–68.
- Mitchell AJ, Sheth B, Gill J, et al. Prevalence and predictors of post-stroke mood disorders: a meta-analysis and meta-regression of depression, anxiety and adjustment disorder. *Gen Hosp Psychiatry*. 2017;47:48–60.
- Garton AL, Sisti JA, Gupta VP, Christophe BR, Connolly ES Jr. Poststroke Post-Traumatic Stress Disorder: A Review [published correction appears in *Stroke*. 2017;48(3):e96]. *Stroke*. 2017;48(2):507–12.
- Ben Assayag E, Tene O, Korczyn AD, et al. Posttraumatic stress symptoms after stroke: the effects of anatomy and coping style. *Stroke*. 2022;53(6):1924–33.
- Rutovic S, Kadojic D, Dikanovic M, Solic K, Malojcic B. Prevalence and correlates of post-traumatic stress disorder after ischaemic stroke. *Acta Neurol Belg*. 2021;121(2):437–42.
- Pedowitz E, Derby L, Cruz GJ, Trainor A, Edmondson D, Cornelius T. Relationship between NIH stroke symptoms and post-traumatic stress disorder in patients evaluated for transient ischemic attack/stroke. *Gen Hosp Psychiatry*. 2021;70:98–102.
- Garton A, Gupta VP, Pucci JU, Couch CK, Connolly ES Jr. Incidence and predictors of post-traumatic stress symptoms in a cohort of patients with intracerebral hemorrhage. *Clin Neurol Neurosurg*. 2020;190: 105657.
- Bruggemann L, Annoni JM, Staub F, von Steinbüchel N, Van der Linden M, Bogousslavsky J. Chronic posttraumatic stress symptoms after nonsevere stroke. *Neurology*. 2006;66(4):513–6.
- Kiphuth IC, Utz KS, Noble AJ, Köhrmann M, Schenk T. Increased prevalence of posttraumatic stress disorder in patients after transient ischemic attack. *Stroke*. 2014;45(11):3360–6.
- Wu S, Wu B, Liu M, et al. Stroke in China: advances and challenges in epidemiology, prevention, and management. *Lancet Neurol*. 2019;18(4):394–405.
- Wang W, Jiang B, Sun H, et al. Prevalence, incidence, and mortality of stroke in China: results from a nationwide population-based survey of 480 687 adults. *Circulation*. 2017;135(8):759–71.
- Kronish IM, Edmondson D, Goldfinger JZ, Fei K, Horowitz CR. Post-traumatic stress disorder and adherence to medications in survivors of strokes and transient ischemic attacks. *Stroke*. 2012;43(8):2192–7.
- Edmondson D, Horowitz CR, Goldfinger JZ, Fei K, Kronish IM. Concerns about medications mediate the association of posttraumatic stress disorder with adherence to medication in stroke survivors. *Br J Health Psychol*. 2013;18(4):799–813.
- Jiang C. Posttraumatic stress disorder after a first-ever intracerebral hemorrhage in the Chinese population: a pilot study. *Appl Neuropsychol Adult*. 2020;27(1):1–8.
- Noble AJ, Baisch S, Mendelow AD, Allen L, Kane P, Schenk T. Posttraumatic stress disorder explains reduced quality of life in subarachnoid hemorrhage patients in both the short and long term. *Neurosurgery*. 2008;63(6):1095–104.
- Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand*. 1983;67(6):361–70.

28. Leung CM, Wing YK, Kwong PK, Lo A, Shum K. Validation of the Chinese-Cantonese version of the hospital anxiety and depression scale and comparison with the Hamilton Rating Scale of Depression. *Acta Psychiatr Scand*. 1999;100(6):456–61.
29. Martin CR, Thompson DR, Chan DS. An examination of the psychometric properties of the Hospital Anxiety and Depression Scale in Chinese patients with acute coronary syndrome. *Psychiatry Res*. 2004;129(3):279–88.
30. Yang Y, Ding R, Hu D, Zhang F, Sheng L. Reliability and validity of a Chinese version of the HADS for screening depression and anxiety in psychocardiological outpatients. *Compr Psychiatry*. 2014;55:215–20.
31. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption—II. *Addiction*. 1993;88(6):791–804.
32. Li Q, Babor TF, Hao W, Chen X. The Chinese translations of Alcohol Use Disorders Identification Test (AUDIT) in China: a systematic review. *Alcohol Alcohol*. 2011;46(4):416–23.
33. Xue C. Alcohol use disorders identification test. *Shanghai Psychiatry*. 1989;1:45,48.[in Chinese].
34. Li B, Sheng Y, Zhang B, et al. The test of AUDIT in China. *Chin Ment Health J*. 2003;17:1–3 [in Chinese].
35. PTSD Checklist for DSM-5 (PCL-5)—PTSD: National Center for PTSD. <https://www.ptsd.va.gov/professional/assessment/adult-sr/ptsd-checklist.asp>. Available from Cited 6 Mar 2020]
36. Sumner JA, Kronish IM, Pietrzak RH, et al. Dimensional structure and correlates of posttraumatic stress symptoms following suspected acute coronary syndrome. *J Affect Disord*. 2015;186:178–85.
37. Cornelius T, Birk JL, Derby L, Ellis J, Edmondson D. Effect of cohabiting partners on the development of posttraumatic stress symptoms after emergency department visits for stroke and transient ischemic attack. *Soc Sci Med*. 2021;281: 114088.
38. Lehman LL, Maletsky K, Beaute J, et al. Prevalence of symptoms of anxiety, depression, and post-traumatic stress disorder in parents and children following pediatric stroke. *J Child Neurol*. 2020;35(7):472–9.
39. Huang RW, Shen T, Ge LM, Cao L, Luo JF, Wu SY. Psychometric properties of the chinese version of the primary care post-traumatic stress disorder screen-5 for medical staff exposed to the covid-19 pandemic. *Psychol Res Behav Manag*. 2021;14:1371–8.
40. Cheng P, Xu LZ, Zheng WH, et al. Psychometric property study of the posttraumatic stress disorder checklist for DSM-5 (PCL-5) in Chinese healthcare workers during the outbreak of corona virus disease 2019. *J Affect Disord*. 2020;277:368–74.
41. Liu P, Wang L, Cao C, et al. The underlying dimensions of DSM-5 posttraumatic stress disorder symptoms in an epidemiological sample of Chinese earthquake survivors. *J Anxiety Disord*. 2014;28(4):345–51.
42. Weathers FW, Blake DD, Schnurr PP, Kaloupek DG, Marx BP, Keane TM. Clinician-Administered PTSD Scale for DSM-5 (CAPS-5). Boston, MA: National Center for PTSD; 2013.
43. Weathers FW, Bovin MJ, Lee DJ, et al. The Clinician-Administered PTSD Scale for DSM-5 (CAPS-5): development and initial psychometric evaluation in military veterans. *Psychol Assess*. 2018;30(3):383–95.
44. Armour C, Tsai J, Durham TA, Charak R, Biehn TL, Elhai JD, et al. Dimensional structure of DSM-5 posttraumatic stress symptoms: Support for a hybrid Anhedonia and Externalizing Behaviors model. *J Psychiatr Res*. 2015;61:106–13.
45. Doll WJ, Xia W, Torkzadeh G. A confirmatory factor analysis of the end-user computing satisfaction instrument. *MIS Q*. 1994;18(4):453–61.
46. Kenny D. Measuring Model Fit. *Davidakenny.Net* (2012) (1985) 1–10. <http://davidakenny.net/cm/fit.htm>. Retrieved from
47. Hu L, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Struct Equ Model Multidiscip J*. 1999;6(1):1–55. <https://doi.org/10.1080/10705519909540118>. Available from.
48. Kraemer HC, Periyakoil VS, Noda A. Kappa coefficients in medical research. *Stat Med*. 2002;21(14):2109–29.
49. Iacobucci D. Structural equations modeling: Fit Indices, sample size, and advanced topics. *J Consum Psychol*. 2010;20:90–8.
50. Liyanage-Don N, Birk J, Cornelius T, et al. Medications as Traumatic Reminders in Patients With Stroke/Transient Ischemic Attack-Induced Posttraumatic Stress Disorder. *Stroke*. 2021;52(1):321–4.
51. Edmondson D, Richardson S, Fausett JK, Falzon L, Howard VJ, Kronish IM. Prevalence of PTSD in survivors of stroke and transient ischemic attack: a meta-analytic review. *PLoS ONE*. 2013;8(6): e66435.
52. van Prooijen J-W, van der Kloot WA. Confirmatory analysis of exploratively obtained factor structures. *Educ Psychol Meas [Internet]*. 2001;61(5):777–92.
53. Svein J, Bondjers K, Willebrand M. Psychometric properties of the PTSD checklist for DSM-5: a pilot study. *Eur J Psychotraumatol*. 2016;7:30165.
54. Wang L, Zhang L, Armour C, et al. Assessing the underlying dimensionality of DSM-5 PTSD symptoms in Chinese adolescents surviving the 2008 Wenchuan earthquake. *J Anxiety Disord*. 2015;31:90–7.
55. Islam MS, Ferdous MZ, Sujon MSH, et al. The psychometric properties of the Bangla Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5): preliminary reports from a large-scale validation study. *BMC Psychiatry*. 2022;22(1):280.
56. Ashbaugh AR, Houle-Johnson S, Herbert C, El-Hage W, Brunet A. Psychometric validation of the english and french versions of the posttraumatic stress disorder checklist for DSM-5 (PCL-5). *PLoS ONE*. 2016;11: e0161645.
57. Wortmann JH, Jordan AH, Weathers FW, Resick PA, Dondanville KA, Hall-Clark B, et al. Psychometric analysis of the PTSD Checklist-5 (PCL-5) among treatment-seeking military service members. *Psychol Assess*. 2016;28:1392–403.
58. Zelazny K, Simms LJ. Confirmatory factor analyses of DSM-5 posttraumatic stress disorder symptoms in psychiatric samples differing in criterion a status. *J Anxiety Disord*. 2015;34:15–23.
59. Ito M, Takebayashi Y, Suzuki Y, Horikoshi M. Posttraumatic stress disorder checklist for DSM-5: psychometric properties in a Japanese population. *J Affect Disord*. 2019;247:11–9.
60. Seligowski AV, Orcutt HK. Support for the 7-factor hybrid model of PTSD in a community sample. *Psychol Trauma*. 2016;8:218–21.
61. Hall BJ, Yip PSY, Garabiles MR, Lao CK, Chan EWW, Marx BP. Psychometric validation of the PTSD checklist-5 among female Filipino migrant workers. *Eur J Psychotraumatol*. 2019;10(1):1571378.
62. Boyd JE, Cameron DH, Shnaider P, McCabe RE, Rowa K. Sensitivity and specificity of the Posttraumatic Stress Disorder Checklist for DSM-5 in a Canadian psychiatric outpatient sample. *J Trauma Stress*. 2022;35(2):424–33.
63. Pereira-Lima K, Loureiro SR, Bolsoni LM, Apolinario da Silva TD, Osório FL. Psychometric properties and diagnostic utility of a Brazilian version of the PCL-5 (complete and abbreviated versions). *Eur J Psychotraumatol*. 2019;10(1):1581020.
64. Yin Q, Sun Z, Liu T, et al. Posttraumatic stress symptoms of health care workers during the corona virus disease 2019. *Clin Psychol Psychother*. 2020;27(3):384–95.
65. Cody MW, Jones JM, Woodward MJ, Simmons CA, Gayle BJ. Correspondence between self-report measures and clinician assessments of psychopathology in female intimate partner violence survivors. *J Interpers Violence*. 2017;32(10):1501–23.
66. Bovin MJ, Marx BP, Weathers FW, et al. Psychometric properties of the PTSD checklist for diagnostic and statistical manual of mental disorders-fifth edition (PCL-5) in veterans. *Psychol Assess*. 2016;28(11):1379–91.
67. Blevins CA, Weathers FW, Davis MT, Witte TK, Domino JL. The posttraumatic stress disorder checklist for DSM-5 (PCL-5): development and initial psychometric evaluation. *J Trauma Stress*. 2015;28(6):489–98.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.