

CORRECTION

Open Access



# Correction: Optimizing psychotherapy dosage for comorbid depression and personality disorders (PsyDos): a pragmatic randomized factorial trial using schema therapy and short-term psychodynamic psychotherapy

Marit Kool<sup>1\*</sup>, Henricus L. Van<sup>1</sup>, Anna Bartak<sup>1</sup>, Saskia C. M. de Maat<sup>1</sup>, Arnoud Arntz<sup>2</sup>, Johanna W. van den Eshof<sup>1</sup>, Jaap Peen<sup>3</sup>, Matthijs Blankers<sup>3,4,5</sup>, Judith E. Bosmans<sup>6</sup> and Jack J. M. Dekker<sup>3,7</sup>

**Correction: BMC Psychiatry 18, 252 (2018)**  
<https://doi.org/10.1186/s12888-018-1829-1>

Following publication of the original article [1], the author identified errors in block randomization and sample size. The correct sentence of block randomization and sample size is given below.

Block randomization

The sentence currently reads:

The original article can be found online at <https://doi.org/10.1186/s12888-018-1829-1>.

\*Correspondence:

Marit Kool

Marit.Kool@arkin.nl

<sup>1</sup> Arkin Mental Health Care, Domselaerstraat 128, 1093 MB Amsterdam, the Netherlands

<sup>2</sup> Department of Clinical Psychology, University of Amsterdam, Amsterdam, the Netherlands

<sup>3</sup> Department of Research, Arkin Mental Health Care, Amsterdam, the Netherlands

<sup>4</sup> Amsterdam UMC, Location AMC, Department of Psychiatry, University of Amsterdam, Amsterdam, the Netherlands

<sup>5</sup> Trimbos Institute – Netherlands Institute of Mental Health and Addiction, Utrecht, the Netherlands

<sup>6</sup> Department of Health Sciences, Faculty of Earth & Life Sciences, Free University Amsterdam, Amsterdam Public Health Research Institute, Amsterdam, The Netherlands

<sup>7</sup> Department of Clinical Psychology, VU University of Amsterdam, Amsterdam, the Netherlands

Patients will then be randomized by one of two research department employees in one of four groups (with a 1:1:1:1 allocation) using a computer script performing block randomization (25 vs 50-sessions; ST vs SPSP).

The sentence should read:

Patients will then be randomized by one of two research department employees in one of four groups (with a 1:1:1:1 allocation) with random allocation sequences that were generated using the SPSS random number generator (SPSS, Chicago).

Sample size

The sentence currently reads:

According to this effect size (and given our choices of  $\alpha = 0.05$ , two-tailed, power  $(1-\beta) = 0.80$ ) 78 patients are needed in both dosage-groups. When 25% dropout is taken into account at least 200 patients will be needed for inclusion.

The sentence should read:

According to this effect size (and given our choices of  $\alpha = 0.05$ , two-tailed, power  $(1-\beta) = 0.80$ ) 79 patients are needed in both dosage-groups. When 25% dropout is taken into account at least 211 patients will be needed for inclusion.

Published online: 16 June 2023



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Reference

1. Kool Marit, et al. Optimizing psychotherapy dosage for comorbid depression and personality disorders (PsyDos): a pragmatic randomized factorial trial using schema therapy and short-term psychodynamic psychotherapy. *BMC Psychiatry*. 2018;18:252. <https://doi.org/10.1186/s12888-018-1829-1>.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

